ERRATUM Open Access



Erratum to: 22q11.2 duplication syndrome: elevated rate of autism spectrum disorder and need for medical screening

Tara L. Wenger^{1,2*†}, Judith S. Miller^{2†}, Lauren M. DePolo², Ashley B. de Marchena², Caitlin C. Clements^{2,3}, Beverly S. Emanuel^{4,5}, Elaine H. Zackai^{4,5,6}, Donna M. McDonald-McGinn^{4,5} and Robert T. Schultz^{2,6}

Erratum

After publication of the research article [1], the authors noticed a minor error in panel **d** of Fig. 3. The authors would like to draw the reader's attention towards the corrected Fig. 3 below. The corrected panel **d** reflects a very minor change which does not change the overall conclusion of the study.

Figure 3: This figure shows the corrected Fig. 3 of the original article.

Author details

¹Department of Pediatrics, Seattle Children's Hospital, M/S OB.9.5204800 Sand Point Way NE, Seattle, WA, USA. ²Center for Autism Research, Children's Hospital of Philadelphia, 3535 Market Street, Philadelphia, PA 19104, USA. ³Department of Psychology, University of Pennsylvania, 3720 Walnut Street, Philadelphia, PA 19104, USA. ⁴Division of Human Genetics and Molecular Biology, Children's Hospital of Philadelphia, 3401 Civic Center Boulevard, Philadelphia, PA 19104, USA. ⁵22q and You Center, Children's Hospital of Philadelphia, 3401 Civic Center Boulevard, Philadelphia, PA 19104, USA. ⁶Department of Pediatrics, University of Pennsylvania, 3401 Civic Center Boulevard, Philadelphia, PA 19104, USA.

Received: 7 July 2016 Accepted: 7 July 2016 Published online: 13 July 2016

References

 Wenger TL, Miller JS, DePolo LM, de Marchena AB, Clements CC, Emanuel BS, et al. 22q11.2 duplication syndrome: elevated rate of autism spectrum disorder and need for medical screening. Mol Autism. 2016;7:27.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- · Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at www.biomedcentral.com/submit



Full list of author information is available at the end of the article



© 2016 The Author(s). **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated.

^{*} Correspondence: Tara.Wenger@seattlechildrens.org †Equal contributors

¹Department of Pediatrics, Seattle Children's Hospital, M/S OB.9.5204800 Sand Point Way NE, Seattle, WA, USA

²Center for Autism Research, Children's Hospital of Philadelphia, 3535 Market Street, Philadelphia, PA 19104, USA




Fig. 3 Questionnaire results for participants with 22q11.2DupS, 22q11.2DupS and comorbid ASD, idiopathic ASD, and typical development. Individuals with idiopathic ASD (*gray*), 22q11.2DS (*red*), and TDC (*green*) were compared to individuals with 22q11.2DupS (*dashed line*) on four parent-report questionnaires about behavioral symptoms. *Error bars* represent one standard error. The 22q11.2DupS group was further divided into individuals who have received a gold standard diagnosis of ASD (DupASD; *light blue*) and those who did not (DupNoASD; *dark blue*). On all measures, individuals with 22q11.2DupS showed scores similar to individuals with 22q11.2Ds. However, when the 22q11.2DupS group was divided into subgroups, individuals in the DupASD subgroup showed scores similar to individuals with idiopathic ASD, whereas individuals in the DupNoASD subgroup showed mean scores in the average ranges, demonstrating less impairment than individuals with 22q11.2Ds. Measures: **a** SRS and **b** CASI-4R: Scores reported in T-scores with mean 50 and SD 10. Scores below 60 considered in normal range. **c** SCQ: Raw scores reported. Scores above 15 are strongly suggestive of ASD. **d** Vineland-II: Scores reported in standard scores with mean 100 and SD 15. Scores above 90 considered in the average range. Abbreviations: ASD, autism spectrum disorder; CASI-4R, Child and Adolescent Symptom Inventory-4R; SCQ, Social Communication Questionnaire; SRS, Social Responsiveness Scale; TDC, typically developing children; Vineland-II, Vineland Adaptive Behavior Scales-II